

Klinische studies naar blauw licht 415nm

Literatuur

- *William Cunliffe J. and Harald Gollnick P.M. (2002). Acne Diagnosis and Management. Martin Dunitz, London.*
- *Marks R. and Plewig G. (1990). Acne and Related Disorders. Martin Dunitz, London.*
- *Kawada A, Aragane Y, Kameyama H, Sangen Y and Tezuka T. (2002). Acne phototherapy with a high-intensity, enhanced, narrowband, blue light source: an open study and in vitro investigation. Journal of Dermatological Science 30.*
- *Elman M, Slatkine M and Harth Y. (2003). The effective treatment of acne vulgaris by a high-intensity, narrow band 405-420nm light source. Journal of Cosmetic & Laser therapy 5.*
- *Omi T, Bjerring P, Sato S, Kawana S, Hankins W.R. and Honda M. (2004). 420 nm intense continuous light therapy for acne. Journal of Cosmetic & Laser therapy 6.*
- *Shalita A.R, Harth Y, Elman M. (2001). Acne PhotoClearing (APC™) using a novel, high-intensity, enhanced, narrow band, blue light source. Clinical Applications Notes, vol.9 No 1.*
- *Lee W.L, Shalita A.R. and Poh-Fritzpatrick M.B. (1978). Comparative studies of Protoporphyrin production in P. acnes and P. granulosum*
- *Arakena K, Ryu A, Hayashi C, Masunaga T, Shinmoto K, Mashiko S, Nagango T and Hirobe M (1996). Singlet oxygen generation from coroporphyrin in P. Acnes on irradiation. Biochemical and biophysical research communications 223 (1996) 579-528.*
- *Koninklijke Nederlandse Maatschappij ter bevordering er Pharmacie (2004). Informatarium Medicamentorum. Haasbeek C., Alphen a/d Rijn.*
- *M. Kruijssen, juni 2005, afstudeerproject Acnetherapie door middel van blauw licht, Hogeschool van Utrecht, Gezondheidszorg*